

## Simple closed geodesics on regular tetrahedra in spaces of constant curvature

Darya Sukhorebska

*B. Verkin Institute for Low Temperature Physics and Engineering of the National Academy of Sciences of Ukraine (B. Verkin ILTPE of NASU)*

suhdaria0109@gmail.com

Alexander Borisenko

*B. Verkin Institute for Low Temperature Physics and Engineering of the National Academy of Sciences of Ukraine (B. Verkin ILTPE of NASU)*

aborisenk@gmail.com

Since regular triangles form a regular tiling of Euclidean plane it easily follows the full classification of closed geodesics on a regular tetrahedron in Euclidean space.

We described all simple (without self-intersections) closed geodesics on regular tetrahedra in three-dimensional hyperbolic and spherical spaces. In these spaces the tetrahedron's curvature is concentrated not only into its vertices but also into its faces. The value  $\alpha$  of the faces' angle for hyperbolic space's tetrahedron satisfies  $0 < \alpha < \pi/3$  and for a tetrahedron in spherical space the faces' angle is measured  $\alpha$  that  $\pi/3 < \alpha < 2\pi/3$ . The intrinsic geometry of such tetrahedra depends on the value of its faces' angles.

A simple closed geodesic on a tetrahedron has the type  $(p, q)$  if it has  $p$  points on each of two opposite edges of the tetrahedron,  $q$  points on each of another two opposite edges, and there are  $(p + q)$  points on each edges of the third pair of opposite one.

We prove that on a regular tetrahedron in hyperbolic space for any coprime integers  $(p, q)$ ,  $0 \leq p < q$ , there exists unique, up to the rigid motion of the tetrahedron, simple closed geodesic of type  $(p, q)$ . These geodesics exhaust all simple closed geodesics on a regular tetrahedron in hyperbolic space. The number of simple closed geodesics of length bounded by  $L$  is asymptotic to constant (depending on  $\alpha$ ) times  $L^2$ , when  $L$  tending to infinity [1].

On a regular tetrahedron in spherical space there exists the finite number of simple closed geodesic. The length of all these geodesics is less than  $2\pi$ . For any coprime integers  $(p, q)$  we presented the numbers  $\alpha_1$  and  $\alpha_2$  depending on  $p, q$  and satisfying the inequalities  $\pi/3 < \alpha_1 < \alpha_2 < 2\pi/3$  such that on a regular tetrahedron in spherical space with the faces' angle of value  $\alpha \in (\pi/3, \alpha_1)$  there exists unique, up to the rigid motion of the tetrahedron, simple closed geodesic of type  $(p, q)$  and on a regular tetrahedron with the faces' angle of value  $\alpha \in (\alpha_2, 2\pi/3)$  there is no simple closed geodesic of type

$(p, q)$ .

## References

- [1] A A Borisenko, D D Sukhorebska, "Simple closed geodesics on regular tetrahedra in hyperbolic space", Mat. Sb., 2020, 211(5), p.3-30. (in Russian). *English translation:* SB MATH, 2020, 211(5), DOI:10.1070/SM9212