

Higher Kazhdan projections, ℓ^2 -Betti numbers & the coarse Baum-Connes conjecture

Sanaz Pooya

based on joint work with Kang Li and Piotr Nowak

Polish Academy of Sciences IMPAN

8th ECM

23 June 2021

Kazhdan's property (T)

In the mid 60's, Kazhdan defined property (T) for locally compact groups and used it as a tool to demonstrate that a large class of lattices in higher rank lie groups are finitely generated. E.g. $SL(n, \mathbb{Z})$ for $n \geq 3$.

Characterisation

A group G has property (T) iff there exists a projection $p \in C_{\max}^* G$ whose image under any unitary representation (π, \mathcal{H}) of G is the orthogonal projection $\mathcal{H} \rightarrow \mathcal{H}^{\pi(G)}$ onto the fixed vectors.

- This projection is called *Kazhdan projection*.
- The Kazhdan projection is unique and non-zero inside $C_{\max}^* G$.
- For an infinite group G , Kazhdan projection in $C_{\text{red}}^* G$ is always zero.
- Existence of this projection violates (surjectivity of) a certain method of proof for the Baum-Connes conjecture, and is a source of counterexamples.

Higher Kazhdan projection

For G finitely generated (hence of type F_1), fix (π, \mathcal{H})

$$p_0: \mathcal{H} \rightarrow \mathcal{H}^{\pi(G)} \quad \text{Kazhdan projection}$$

\rightsquigarrow higher degrees, one may use the identification $\mathcal{H}^{\pi(G)} = H^0(G, \mathcal{H})$

Definition (Li-Nowak-P 2020)

Let G be a discrete group of type F_{n+1} . Let (π, \mathcal{H}) be a unitary representation of G . The higher Kazhdan projection in degree n associated with π is the orthogonal projection

$$p_n: \mathcal{H}^{\oplus k_n} \rightarrow \tilde{H}^n(G, \mathcal{H})$$

Remark

- It always lives in the matrices over the von Neumann algebra generated by $\pi(G)$.
- Assuming spectral gap for the higher Laplacian $\pi(\Delta_n)$, it lives in the matrices over the C^* -algebra generated by $\pi(G)$.

ℓ^2 -Betti numbers

group von Neumann algebra $LG = \overline{\mathbb{C}G}^{SOT} \subseteq \mathcal{B}(\ell^2 G)$

ℓ^2 -Betti number

$$\beta_{(2)}^n(G) = \dim_{LG} \tilde{H}^n(G, \ell^2 G) \in [0, \infty]$$

canonical trace $\tau: LG \rightarrow \mathbb{C}$ defined by $\tau(\sum_{\text{finite}} c_g g) = c_e$

$$\tilde{H}^n(G, \ell^2 G) = p_n(\ell^2 G^{\oplus k_n}) \quad \text{right } LG\text{-module}$$

$$\beta_{(2)}^n(G) = \dim_{LG} \tilde{H}^n(G, \ell^2 G) = \dim_{LG} p_n(\ell^2 G^{\oplus k_n}) = (\text{Tr} \otimes \tau)(p_n)$$

Identifying higher Kazhdan projections

Proposition (Folklore)

Assume $\lambda(\Delta_n)$ has spectral gap so that p_n belongs to $M_{k_n}(C_{red}^*G)$.
Then we have that

$$\beta_{(2)}^n(G) = \tau_*([p_n])$$

In particular

- if $[p_n] = 0$ in $K_0(C_{red}^*G)$, then $\beta_{(2)}^n(G) = 0$
- if $[p_n] \in \mathbb{Z} \cdot [1]$, then $\beta_{(2)}^n(G) \in \mathbb{Z}$

Example

- $K_0(C_{red}^*\mathbb{F}_n) = \mathbb{Z} \cdot [1]$, and $\beta_{(2)}^1(\mathbb{F}_n) = n - 1 \rightsquigarrow [p_1] = (n - 1)[1]$
- $\beta_{(2)}^1(PSL(2, \mathbb{Z})) = 1/6 \rightsquigarrow [p_1] \notin \mathbb{Z} \cdot [1]$

Coarse Baum-Connes conjecture

X discrete metric space with bounded geometry

\mathcal{H} separable, infinite dimensional Hilbert space

$\mathbb{C}[X] \subseteq \mathcal{B}(\ell^2(X, \mathcal{H}))$: $*$ -algebra of finite propagation operators with compact entries $T_{(x,y)}$.

Roe algebra $C^*[X] = \overline{\mathbb{C}[X]} \subseteq \mathcal{B}(\ell^2(X, \mathcal{H}))$

Coarse Baum-Connes conjecture, Roe, 1993

For all X with bounded geometry the coarse assembly map

$$\mu_{\bullet} : KK_{\bullet}(X) \rightarrow K_{\bullet}(C^*[X]) \quad \bullet = 0, 1$$

is an isomorphism.

Application to the coarse Baum-Connes conjecture

- $\beta^n(G) = \dim_{\mathbb{C}} H^n(G, \mathbb{C}) \in \mathbb{N}$
- $\beta_{(2)}^n(G) = \dim_{L G} \tilde{H}^n(G, \ell^2 G) \in [0, \infty]$

Theorem (Li-Nowak-P 2020)

Let G be an exact residually finite group of type F_{n+1} . Let $N = \{N_i\}_i$ be a filtration of finite index normal subgroups of G . Let $\pi = \bigoplus_i \lambda_i$. Assume that $\pi(\Delta_n)$ has a spectral gap such that p_n belongs to $M_{k_n}(\mathbb{C}_N^* G)$. If the coarse Baum-Connes assembly map for the box space $Y = \coprod G/N_i$ of G is surjective, then

$$\beta_{(2)}^n(N_i) = \beta^n(N_i)$$

for sufficiently large i .

- ↪ strategy to find counterexamples to the conjecture
- ↪ consequences of surjectivity of the conjecture

Thanks for listening!